



Available online at [www.sciencedirect.com](http://www.sciencedirect.com)

ScienceDirect

journal homepage: [www.elsevier.com/locate/psyneuen](http://www.elsevier.com/locate/psyneuen)



# Psychological and cortisol reactivity to experimentally induced stress in adults with ADHD

Sivan Raz<sup>a,b,\*</sup>, Dmitry Leykin<sup>a,c</sup>

<sup>a</sup> Department of Psychology, Tel Hai College, 12208, Israel

<sup>b</sup> Departments of Behavioral Sciences and Psychology, The Center for Psychobiological Research, The Max Stern Yezreel Valley College, 19300, Israel

<sup>c</sup> Recanati School for Community Health Professions, Department of Emergency Medicine, Faculty of Health Sciences, Ben-Gurion University of the Negev, Beer-Sheva, Israel

Received 28 January 2015; received in revised form 4 May 2015; accepted 4 May 2015

## KEYWORDS

ADHD;  
Adults;  
Test anxiety;  
Psychological stress;  
Cortisol reactivity;  
HPA axis

**Summary** Individuals with ADHD suffer from increased vulnerability to environmental and mental stressors and may be at increased risk for chronic stress in everyday life. The Hypothalamic-Pituitary-Adrenal (HPA) axis is a critical physiological system that mediates responses to stress. The present study seeks to examine test performance, test anxiety, self-reported psychological stress and cortisol reactivity to mental-cognitive stress in adults with ADHD when compared with healthy controls. Stress was induced by an arithmetic ability test. Psychological stress was assessed repeatedly throughout the experimental session. Salivary cortisol, an indicator of the HPA axis function, was evaluated immediately upon arrival, as well as 1 min and 20 min post-test completion. Results revealed higher levels of test anxiety and poorer performance on the test in the ADHD group. The ADHD and control groups showed no difference in base-line levels of subjective stress and in subjective stress levels 20 min after the test. In contrast, individuals with ADHD reported significantly higher levels of stress at the test anticipation phase and 1 min post-test completion. Cortisol response to stress differed according to group: in the ADHD group, 20 min post-test cortisol levels were significantly higher than base-line cortisol levels. This was not evident in the control group. These results suggest greater activation of the HPA axis in response to stress in adults with ADHD when compared with healthy controls. Adults with ADHD do not differ from controls in basal levels of subjective

\* Corresponding author at: Departments of Behavioral Sciences and Psychology, The Center for Psychobiological Research, The Max Stern Yezreel Valley College, 19300, Emek Yezreel, Israel. Tel.: +972 54 6634301; fax: +972 46423618.

E-mail address: [sivanr@yvc.ac.il](mailto:sivanr@yvc.ac.il) (S. Raz).



Corominas et al. (2013) examined CAR in adults with ADHD in order to assess possible differences between the combined and inattentive ADHD subtypes. Results showed no significant differences in the mean increase of CAR between the inattentive and combined subtypes. This study did not include a comparison group of healthy individuals. Hirvikoski et al. (2009) compared adults with ADHD with healthy controls regarding diurnal salivary cortisol in the everyday environment and salivary cortisol before and after cognitive stress in a laboratory setting. Results of this study showed that individuals with ADHD reported significantly more self-perceived stress than controls, and subjective stress correlated with the amount of stressors in everyday life. The two groups were comparable with respect to overall diurnal cortisol levels and rhythm, as well as in pre- and post-stress cortisol concentrations. Post-stress cortisol (but not baseline cortisol) concentration was positively correlated with impulsivity. The group with high post-stress cortisol also reported higher levels of self-perceived stress and more stressors in everyday life. The diagnosis of ADHD significantly increased the risk of belonging to the group with high post-stress cortisol levels. However, this study was limited by a high incidence of comorbid disorders, since participants were recruited from a Neuropsychiatric Unit of a University Hospital, with 57% of the ADHD group having at least one additional diagnosis (e.g., depression, panic disorder, anxiety and borderline personality). Lackschewitz et al. (2008) examined psychological and physiological stress responses in adult ADHD subjects in comparison with healthy controls under laboratory conditions. Participants with ADHD reported greater subjective stress but no group differences in cortisol reactivity were seen. The sample size in this study was relatively small ( $n = 18$ ) and one-third of the ADHD group had comorbid psychiatric disorders.

Young adults with ADHD often demonstrate academic difficulties that result in significantly lower achievement scores and less overall academic success than peers (Barkley, 2006; Barkley et al., 2008; Frazier et al., 2007; Weyandt and DuPaul, 2006). One factor that has been strongly suggested as a contributor to lower academic performance, in the general population, is test-related anxiety (Chapell et al., 2005; Zeidner, 1998). Reaser et al. (2007) found that college students with ADHD tend to possess a negative attribution style with regard to their test performance as well as motivational deficiencies. Nevertheless, very few studies have directly assessed levels of test anxiety in adults with ADHD (Dan and Raz, 2012; Lewandowski et al., 2013; Nelson et al., 2014). Test anxiety may be associated with a more negative self-perception as well as with poorer test performance, greater self-perceived stress and alterations in cortisol reactivity to psychological stress in individuals with ADHD.

The aim of the present study is to examine test anxiety, test performance, self-reported psychological stress and cortisol reactivity to mental-cognitive stress (arithmetic ability test) in adults with ADHD (without comorbid psychiatric conditions), in comparison with healthy controls. We expected participants with ADHD to report higher basal (trait) levels of test anxiety and to demonstrate poorer performance (lower grades) on the arithmetic test. Based on the relevant literature, it was hypothesized that ADHD adults and healthy controls would show comparable baseline levels of both subjective stress and cortisol prior to exposure to the

stress manipulation. However, in response to the laboratory stressor, higher levels of subjective stress accompanied by alterations in cortisol reactivity in the ADHD group compared with controls were expected.

## 2. Methods and materials

### 2.1. Participants

Forty nine undergraduate students (mean age  $26.62 \pm 1.88$  years) participated in this study. They were divided into two groups: participants diagnosed with ADHD ( $n = 24$ ; 16 females, mean age  $27.16 \pm 1.83$ ), and healthy controls without ADHD ( $n = 25$ ; 19 females, mean age  $26.04 \pm 1.80$ ). Inclusion criteria for the ADHD group were (a) at least six symptoms either of inattention or of hyperactivity-impulsivity on the DSM-IV based ADHD Rating Scale (DuPaul et al., 1998), and (b) previous professional diagnosis: participants provided documentation of valid professional diagnosis of adult ADHD previously made by either a neurologist or psychiatrist from an established clinic in the field of psychoeducational assessment (the formal assessment must include a sound clinical interview). In many cases these documents were the ones submitted to the College's Support Center for Students with Learning Disabilities in order to get special exam accommodations. Students who could not provide such relevant documents were not included in the study.

Inclusion criteria for the control group were (a) fewer than four symptoms either of inattention or of hyperactivity-impulsivity on the ADHD Rating Scale-IV, and (b) no previous diagnosis of ADHD. The ADHD and control groups did not differ with respect to mean age, education or ethnicity distribution. Participants with a prior history or comorbidity with other neurological or mental/psychiatric disorders were excluded from the study. Participants were asked several questions to control for past or present mental/psychiatric disorders, e.g., "Are you suffering, or have you ever been diagnosed as suffering from some kind of mental/psychiatric/personality disorder?", "Are you taking, or have you ever been treated with, some kind of psychiatric medication?" Participants were recruited using departmental mailing lists and through snowball sampling. Written informed consent was obtained from all participants and they were given course credit according to their academic requirements. The study was approved by the Institutional Review Board and is in accordance with the declaration of Helsinki.

### 2.2. Measures

#### 2.2.1. ADHD questionnaire

The assessment questionnaire for ADHD (DuPaul et al., 1998) included 18 items based on the symptoms listed in the DSM-IV for ADHD diagnosis. These symptoms include measurements of attentiveness (9 items), hyperactivity and impulsivity (9 items). Participants were asked to choose whether each described situation was correct or incorrect








post-test (Cort3) than base-line levels (Cort1) and immediately post-test levels (Cort2) ( $p_s=0.0001$ ) (Figure 4a). Another way to look at these results is to calculate the difference in cortisol levels between Cort3 and Cort1 and between Cort3 and Cort2 (Cort3–Cort1; Cort3–Cort2), and to compare those differences between the ADHD and control groups. *T*-test analysis revealed that the difference in cortisol between Cort3 and Cort1 was significantly higher in the


ADHD group than in the control group [ $t(47)=2.16$ ,  $p=0.036$ , Cohen's  $d=0.65$ ]. Similar results were found for the difference in cortisol between Cort3 and Cort2 [ $t(47)=3.39$ ,  $p=0.001$ , Cohen's  $d=1.00$ ] (Figure 4b). These results suggest that the cortisol response to the test was significantly more pronounced in participants with ADHD than in controls.

#### 4. Discussion

In the present study we examined test anxiety, test performance, psychological stress and cortisol reactivity to stress in adults with ADHD compared with healthy controls. We found that participants with ADHD had significantly higher base line levels of Cognitive Obstruction test anxiety, Tenseness test anxiety and Total test anxiety. It seems that individuals with ADHD tend to perceive test situations as more threatening than those without ADHD. Anxiety disorders are a common comorbid condition among individuals with ADHD, with prevalence commonly ranging from 15% to 35%, and even reaching as much as 50% (Kessler et al., 2006; Mancini et al., 1999; Schatz and Rostain, 2006). Despite the well documented comorbidity of ADHD with anxiety disorders, very few studies have directly evaluated test anxiety among adults with ADHD (Dan and Raz, 2012; Lewandowski et al., 2013; Nelson et al., 2014). The current results are in line with Nelson et al. (2014) who found that college students with ADHD reported higher total test anxiety as well as specific aspects of test anxiety, including worry (i.e., cognitive aspects of test anxiety) and emotionality (i.e., physiological aspects of test anxiety). Lewandowski et al. (2013) reported significant differences between ADHD and non-ADHD students regarding their perceptions of, and anxiety during, test taking. Those with ADHD perceived themselves as having more difficulty in reading under timed conditions and reported more test-related anxiety than their peers. Adults with ADHD often demonstrate academic difficulties that result in lower grades than peers and less academic success overall (Barkley, 2006; Barkley et al., 2008; DuPaul and Volpe, 2009; Frazier et al., 2007). Test anxiety may be a major factor contributing to such lower academic performance; indeed, the results of the present study show the performance of participants with ADHD to be significantly poorer on the arithmetic test. Worry, which is considered a fundamental cognitive component of test anxiety, has been found to be particularly related to academic under-achievement (Wine, 1971; Wong, 2008; Zeidner, 1998). The Worry component closely corresponds to the Cognitive Obstruction scale of the test anxiety questionnaire used in this study. Attentional interpretation of the Worry component that may be specifically relevant to ADHD has been suggested by several researchers (Sarason, 1984; Tobias, 1985; Wine, 1971). According to this interpretation, Worry is an attention-demanding cognitive activity involving self-preoccupying, intrusive thoughts that interfere with task-focused thinking. Thus, high test-anxious individuals may be more susceptible to cognitive interference, deficits in allocation of attention and reduction in the cognitive capacity for task solution when confronted with challenging tasks; this may prove to be even truer for individuals with ADHD. It is difficult to determine whether students with ADHD perform less successfully due to test anxiety or



**Figure 3** Differences between ADHD and control groups in levels of self-reported stress on the Visual Analog Scale (VAS). Error bars represent SEM. Asterisks are showing the results of a comparison between groups: \* $p < 0.05$ , \*\* $p < 0.01$ .



**Figure 4** (a) Salivary cortisol in ADHD and control groups at base-line (Cort1), immediately after the test (Cort2) and 20 min after the test (Cort3). (b) Differences in cortisol between Cort3 and Cort1 and between Cort3 and Cort2 for the ADHD and control groups. Error bars represent SEM. Asterisks are showing the results of a comparison between groups: \* $p < 0.05$ , \*\* $p < 0.01$ .







- problems: the impact of callous unemotional traits. *Psychiatry Res.* 187, 204–209.
- Tobias, S., 1985. Test anxiety: interference, defective skills, and cognitive capacity. *Educ. Psychol.* 20, 135–142.
- van West, D., Claes, S., Deboutte, D., 2009. Differences in hypothalamic-pituitary-adrenal axis functioning among children with ADHD predominantly inattentive and combined types. *Eur. Child Adolesc. Psychiatry* 18, 543–553.
- von Dawans, B., Kirschbaum, C., Heinrichs, M., 2011. The Trier Social Stress Test for Groups (TSST-G): a new research tool for controlled simultaneous social stress exposure in a group format. *Psychoneuroendocrinology* 36, 514–522.
- Weyandt, L.L., DuPaul, G.J., 2006. ADHD in college students: a review of the literature. *J. Atten. Dis.* 10, 9–19.
- Willcutt, E.G., 2012. The prevalence of DSM-IV attention-deficit/hyperactivity disorder: a meta-analytic review. *Neurotherapeutics* 9, 490–499.
- Wine, J., 1971. Test anxiety and direction of attention. *Psychol. Bull.* 76, 92–104.
- Wolraich, M.L., McKeown, R.E., Visser, S.N., Bard, D., Cuffe, S., Neas, B., 2012. The prevalence of ADHD: Its diagnosis and treatment in four school districts across two states. *J. Atten. Dis.*, <http://dx.doi.org/10.1177/1087054712453169>.
- Wong, S.S., 2008. The relations of cognitive triad, dysfunctional attitudes, automatic thoughts, and irrational beliefs with test anxiety. *Curr. Psychol.* 27, 177–191.
- Yim, I.S., Quas, J.A., Cahill, L., Hayakawa, C.M., 2010. Children's and adults' salivary cortisol responses to an identical psychosocial laboratory stressor. *Arch. Clin. Neuropsychol.* 20, 727–744.
- Zeidner, M., 1998. *Test Anxiety: The State of the Art*. Plenum, New York.